

CLASSIFICATION CONFIDENTIAL

CONFIDENTIAL

CENTRAL INTELLIGENCE AGENCY  
INFORMATION FROM  
FOREIGN DOCUMENTS OR RADIO BROADCASTS

REPORT

50X1-HUM

CD NO.

COUNTRY USSR

DATE OF  
INFORMATION 1950SUBJECT Scientific - Physics, plastic deformation  
of beams

DATE DIST. 71 Nov 1950

HOW  
PUBLISHED Thrice-monthly periodicalWHERE  
PUBLISHED Moscow

NO. OF PAGES 2

DATE  
PUBLISHED 21 Jul 1950

LANGUAGE Russian

SUPPLEMENT TO  
REPORT NO.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE  
OF THE UNITED STATES WITHIN THE MEANINGS OF ESPIONAGE ACT 50  
U. S. C. 31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION  
OF ITS CONTENTS IN ANY MANNER TO AN UNAUTHORIZED PERSON IS PRO-  
HIBITED BY LAW. REPRODUCTION OF THIS FORM IS PROHIBITED.

THIS IS UNEVALUATED INFORMATION

SOURCE Doklady Akademii Nauk SSSR, Vol LXXIII, No 3, 1950, pp 471-474.

APPLIED THEORY OF PLASTIC  
DEFORMATION OF BEAMS

Yu. I. Yagn and Ye. N. Tarasenko  
Submitted by Acad Smirnov  
29 May 1950

[In this report, of which the following is a digest, an impor-  
tant nonlinear differential equation is employed in an actual ap-  
plied case of plastic deformation, and its solution is indicated  
by current methods.]

In G. Yu. Dzhaneldidze's work (1) the applied theories of elastic deforma-  
tion of beams were widely generalized on the base of essential kinematic schemes  
that take into account the effect of warping of cross sections caused by twist-  
ing. He proposed, in a study of deformation beyond the elastic limit, a simpli-  
fied kinematic model constructed on the assumption that the cross sections re-  
main plane during all forms of deformation of a beam, including torsion.

However, experiments clearly reveal a strong warping of the cross sections  
during all stages of plastic torsion. In this case the observed forms and dimen-  
sions of the warping of cross sections approach those obtained for the same degree  
of torsion in the elastic stage of deformation.

Thus, we are compelled to recommend that in any construction of an applied  
theory of plastic deformation of beams one must preserve the kinematic scheme  
that applies, or responds, to the elastic stage, as was also advocated by V. V.  
Novozhilov (2).

When these recommendations are followed, we find the general equation de-  
scribing the kinematic picture of deformation to be the following nonlinear one:

$$\tau \tau^2 + A \tau^2 + B \tau + C \tau^2 \tau + D \tau^3 + E \tau + G = 0$$

where tau  $\tau$  is the relative angle of twist and A, B, C, D, and E are constants.  
This equation can be approximately solved by B. G. Galerkin's variational method  
or by numerical integration.

- 1 -

CONFIDENTIAL

CLASSIFICATION		CONFIDENTIAL		DISTRIBUTION									
STATE	<input checked="" type="checkbox"/>	NAVY	<input checked="" type="checkbox"/>	NSRB									
ARMY	<input checked="" type="checkbox"/>	AIR	<input checked="" type="checkbox"/>	FBI									

CONFIDENTIAL

CONFIDENTIAL

50X1-HUM

This equation results from the simultaneous solution of the following set:

$$\begin{aligned}\varepsilon_z &= \varepsilon + K_1 y - K_2 x + \tau \varphi \\ \gamma_{xz} &= \tau \left( \frac{\partial \varphi}{\partial x} - y \right) + \bar{\gamma}_{xz} \\ \gamma_{yz} &= \tau \left( \frac{\partial \varphi}{\partial y} + x \right) + \bar{\gamma}_{yz}\end{aligned}$$

where  $\varepsilon_z$  is the longitudinal relative elongation;  $\gamma_{xz}$  and  $\gamma_{yz}$  are the angular displacements in the longitudinal planes parallel to the principal central axes of inertia of the cross section;  $\varepsilon$  is the average value of  $\varepsilon_z$  over the cross section;  $K_1$  and  $K_2$  are the components of the curvature of the beam's axis in the planes of the principal central axes of inertia;  $x$  and  $y$  are the coordinates of this system;  $\varphi$  is a function of torsion satisfying the equation  $\Delta \varphi = 0$  and the boundary conditions  $\partial \varphi / \partial n = y \cos(n, x) - x \cos(n, y)$

and  $\int \varphi d\Omega$  for the determination of the constant;  $\bar{\gamma}_{xz}$  and  $\bar{\gamma}_{yz}$  are the additional angular displacements over and above those displacements that are due to pure torsion. The presence of these terms distinguish our equations from the corresponding ones used by G. Yu. Dzhanelidze for the elastic stage.

[The remainder of the article reduces this set to the nonlinear equation, after suitable transformations and eliminations, with approximations and neglect of high-order terms; A, B, C, D, E are lumped constants.]

## BIBLIOGRAPHY

1. G. Yu. Dzhanelidze. Author's Report of Dissertation: The Statics of Elastic-Plastic Beams (Avtoreferat dissertatsii Statika uprugoplasticheskikh sterzhney). Leningrad, 1949. Dok Ak Nauk SSSR, 66, No 4, 1949.
2. V. V. Novozhilov. Fundamentals of the Nonlinear Theory of Elasticity (Osnovy nelineynoy teorii uprugosti), Section 49, 1948.

- E N D -

- 2 -

CONFIDENTIAL

CONFIDENTIAL